



Entrez PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

Sgo

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Caddy

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Protein Sci. 1999 Dec;8(12):2589-97.[Related Articles, Links](#)

The structures of the neurotrophin 4 homodimer and the brain-derived neurotrophic factor/neurotrophin 4 heterodimer reveal a common Trk-binding site.

Robinson RC, Radziejewski C, Spraggon G, Greenwald J, Kostura MR, Burtnick LD, Stuart DI, Choe S, Jones EY.

Structural Biology Laboratory, The Salk Institute, La Jolla, California 92037, USA.
bob@sbl.salk.edu

The neurotrophins are growth factors that are involved in the development and survival of neurons. Neurotrophin release by a target tissue results in neuron growth along the neurotrophin concentration gradient, culminating in the eventual innervation of the target tissue. These activities are mediated through trk cell surface receptors. We have determined the structures of the heterodimer formed between brain-derived neurotrophic factor (BDNF) and neurotrophin 4 (NT4), as well as the structure of homodimer of NT4. We also present the structure of the Neurotrophin 3 homodimer, which is refined to higher resolution than previously published. These structures provide the first views of the architecture of the NT4 protomer. Comparison of the surface of a model of the BDNF homodimer with the structure of the neurotrophin homodimers reveals common features that may be important in the binding between the neurotrophins and their receptors. In particular, there exists an analogous region on the surface of each neurotrophin that is likely to be involved in trk receptor binding. Variations in sequence on the periphery of this common region serve to confer trk receptor specificity.

PMID: 10631974 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Nov 21 2003 06:56:44